

Bibliography

- [1] M. H. Anderson, J. R. Ensher, M. R. Matthews, C. E. Wieman, and E. A. Cornell, "Observation of Bose-Einstein condensation in a dilute atomic vapor," *Science* **269**, 198–201 (1995).
- [2] C. S. Wood, S. C. Bennett, D. Cho, B. P. Masterson, J. L. Roberts, C. E. Tanner, and C. E. Wieman, "Measurement of Parity Nonconservation and an Anapole Moment in Cesium," *Science* **275**, 1759–63 (1997).
- [3] G. Gwinner, J. A. Behr, S. B. Cahn, A. Ghosh, L. A. Orozco, G. D. Sprouse, and F. Xu, "Magneto-optic trapping of radioactive ^{79}Rb ," *Phys. Rev. Lett.* **72**, 3795–8 (1994).
- [4] O. Stern and W. Gerlach, "Der experimentelle Nachweis des magnetischen Moments des Silberatoms," *Z. Phys.* **8**, 110–1 (1921).
- [5] W. Gerlach and O. Stern, "Der experimentelle Nachweis der Richtungsquantelung im Magnetfeld," *Z. Phys.* **9**, 349–52 (1922).
- [6] N. F. Ramsey, Molecular Beams (Clarendon Press, Oxford, 1956).
- [7] W. D. Phillips and H. Metcalf, "Laser deceleration of an atomic beam," *Phys. Rev. Lett.* **48**, 596–9 (1982).
- [8] W. Ertmer, R. Blatt, J. L. Hall, and M. Zhu, "Laser manipulation of atomic beam velocities: demonstration of stopped atoms and velocity reversal," *Phys. Rev. Lett.* **54**, 996 – 9 (1985).
- [9] S. Chu, L. Hollberg, J. E. Bjorkholm, A. Cable, and A. Ashkin, "Three-dimensional viscous confinement and cooling of atoms by resonant radiation pressure," *Phys. Rev. Lett.* **55**, 48–51 (1985).
- [10] J. P. Gordon and A. Ashkin, "Motion of atoms in a radiation trap," *Phys. Rev. A* **21**, 1606–17 (1980).
- [11] P. D. Lett, R. N. Watts, C. I. Westbrook, W. D. Phillips, P. L. Gould, and H. J. Metcalf, "Observation of Atoms Laser Cooled below the Doppler Limit," *Phys. Rev. Lett.* **61**, 169–72 (1988).

- [12] J. Dalibard and C. Cohen-Tannoudji, “Laser cooling below the Doppler limit by polarization gradients: simple theoretical models,” *J. Opt. Soc. Am. B* **6**, 2023–45 (1989).
- [13] P. J. Ungar, D. S. Weiss, E. Riis, and S. Chu, “Optical molasses and multilevel atoms: theory,” *J. Opt. Soc. Am. B* **6**, 2058–71 (1989).
- [14] E. L. Raab, M. Prentiss, A. Cable, S. Chu, and D. E. Pritchard, “Trapping of neutral sodium atoms with radiation pressure,” *Phys. Rev. Lett.* **59**, 2631–4 (1987).
- [15] C. Monroe, W. Swann, H. Robinson, and C. Wieman, “Very cold trapped atoms in a vapor cell,” *Phys. Rev. Lett.* **65**, 1571 – 4 (1990).
- [16] S. Bali, K. M. O’Hara, M. E. Gehm, S. R. Granade, and J. E. Thomas, “Quantum-diffractive background gas collisions in atom-trap heating and loss,” *Phys. Rev. A* **60**, R29 – 32 (1999).
- [17] H. C. W. Beijerinck, “Rigorous calculation of heating in alkali traps by background gas collisions,” submitted to *Phys. Rev. A*.
- [18] C. J. Myatt, N. R. Newbury, R. W. Ghrist, L. Loutzenhiser, and C. E. Wieman, “Multiply loaded magneto-optical trap,” *Opt. Lett.* **21**, 290 – 2 (1996).
- [19] C. J. Myatt, E. A. Burt, R. W. Ghrist, E. A. Cornell, and C. E. Wieman, “Production of two overlapping Bose-Einstein condensates by sympathetic cooling,” *Phys. Rev. Lett.* **78**, 586–9 (1997).
- [20] E. Riis, D. S. Weiss, K. A. Moler, and S. Chu, “Atom funnel for the production of a slow, high-density atomic beam,” *Phys. Rev. Lett.* **64**, 1658–61 (1990).
- [21] J. Nellesen, J. Werner, and W. Ertmer, “Magneto-optical compression of a monoenergetic sodium atomic beam,” *Opt. Comm.* **78**, 300–308 (1990).
- [22] A. Scholz, M. Christ, D. Doll, J. Ludwig, and W. Ertmer, “Magneto-optical preparation of a slow, cold and bright Ne* atomic beam,” *Opt. Comm.* **111**, 155–62 (1994).
- [23] J. Yu, J. Djemaa, P. Nosbaum, and P. Pillet, “Funnel with orientated Cs atoms,” *Opt. Comm.* **112**, 136–40 (1994).
- [24] Z.-T. Lu, K. L. Corwin, M. J. Renn, M. H. Anderson, E. A. Cornell, and C. E. Wieman, “Low-Velocity Intense Source of Atoms from a Magneto-optical Trap,” *Phys. Rev. Lett.* **77**, 3331–4 (1996).
- [25] C. Y. Park, M. S. Jun, and D. Cho, “Magneto-optical trap loaded from a low-velocity intense source,” *J. Opt. Soc. Am. B* **16**, 994–7 (1999).
- [26] R. S. Williamson III, P. A. Voytas, R. T. Newell, and T. Walker, “A magneto-optical trap loaded from a pyramidal funnel,” *Opt. Express* **3**, 111–117 (1998).

- [27] K. I. Lee, J. A. Kim, H. R. Noh, and W. Jhe, “Single-beam atom trap in a pyramidal and conical hollow mirror,” *Opt. Lett.* **21**, 1177–9 (1996).
- [28] D. Müller, E. A. Cornell, D. Z. Anderson, and E. R. I. Abraham, “Guiding laser-cooled atoms in hollow core fibers,” submitted .
- [29] D. Müller, D. Z. Anderson, R. J. Grow, P. D. D. Schwindt, and E. A. Cornell, “Guiding atoms around curves with lithographically patterned current-carrying wires,” *Phys. Rev. A* (accepted) (1999).
- [30] M. Stephens and C. Wieman, “High collection efficiency in a laser trap,” *Phys. Rev. Lett.* **72**, 3787 – 90 (1994).
- [31] M. Stephens, R. Rhodes, and C. Wieman, “Study of wall coatings for vapor-cell laser traps,” *J. Appl. Phys.* **76**, 3479 – 88 (1994).
- [32] K. Lindquist, M. Stephens, and C. Wieman, “Experimental and theoretical study of the vapor-cell Zeeman optical trap,” *Phys. Rev. A* **46**, 4082 – 90 (1992).
- [33] T. Dinneen, A. Ghiorso, and H. Gould, “An orthotropic source of thermal atoms,” *Rev. Sci. Inst.* **67**, 752 – 5 (1996).
- [34] Z.-T. Lu, K. L. Corwin, K. R. Vogel, C. E. Wieman, T. P. Dinneen, J. Maddi, and H. Gould, “Efficient Collection of ^{221}Fr into a Vapor Cell Magneto-optical Trap,” *Phys. Rev. Lett.* **79**, 994 – 997 (1997).
- [35] J. S. Grossman, L. A. Orozco, M. R. Pearson, J. E. Simsarian, G. D. Sprouse, and W. Z. Zhao, “Hyperfine Anomaly Measurements in Francium Isotopes and the Radial Distribution of Neutrons,” *Phys. Rev. Lett.* **83**, 935–8 (1999).
- [36] M. A. Rowe, S. J. Freedman, B. K. Fujikawa, G. Gwinner, and S.-Q. Shang, “Ground-state hyperfine measurement in laser-trapped radioactive ^{21}Na ,” *Phys. Rev. A* **59**, 1869–73 (1999).
- [37] J. E. Bjorkholm, R. R. Freeman, A. Ashkin, and D. B. Pearson, “Observation of Focusing of Neutral Atoms by the Dipole Forces of Resonance-Radiation Pressure,” *Phys. Rev. Lett.* **41**, 1361–64 (1978).
- [38] S. Chu, J. E. Bjorkholm, A. Ashkin, and A. Cable, “Experimental observation of optically trapped atoms,” *Phys. Rev. Lett.* **57**, 314–7 (1986).
- [39] J. D. Miller, R. A. Cline, and D. J. Heinzen, “Far-off-resonance optical trapping of atoms,” *Phys. Rev. A* **47**, R4567–70 (1993).
- [40] R. A. Cline, J. D. Miller, M. R. Matthews, and D. J. Heinzen, “Spin relaxation of optically trapped atoms by light scattering,” *Opt. Lett.* **19**, 207–209 (1994).

- [41] K. L. Corwin, S. J. M. Kuppens, D. Cho, and C. E. Wieman, “Spin-Polarized Atoms in a Circularly Polarized Optical Dipole Trap,” *Phys. Rev. Lett.* **83**, 1311–14 (1999).
- [42] C. Cohen-Tannoudji and J. Dupont-Roc, “Experimental study of Zeeman light shifts in weak magnetic fields,” *Phys. Rev. A* **5**, 968–84 (1972).
- [43] D. Cho, “Analogous Zeeman effect from the tensor polarizability in alkali atoms,” *J. of Korean Phys. Soc.* **30**, 373–6 (1997).
- [44] S. J. M. Kuppens, K. L. Corwin, K. Miller, T. Chupp, and C. E. Wieman, “The physics of loading an optical dipole trap,” submitted (1999).
- [45] N. R. Newbury, C. J. Myatt, E. A. Cornell, and C. E. Wieman, “Gravitational sisyphus cooling of ^{87}Rb in a magnetic trap,” *Phys. Rev. Lett.* **74**, 2196–9 (1995).
- [46] K. Gibble and S. Chu, “Laser-cooled Cs frequency standard and a new measurement of the frequency shift due to ultracold collisions,” *Phys. Rev. Lett.* **70**, 1771–4 (1993).
- [47] D. W. Keith, C. R. Ekstrom, Q. A. Turchette, and D. E. Pritchard, “An interferometer for atoms,” *Phys. Rev. Lett.* **66**, 2693–6 (1991).
- [48] M. J. Renn, E. A. Donley, E. A. Cornell, C. E. Wieman, and D. Z. Anderson, “Evanescent-wave guiding of atoms in hollow optical fibers,” *Phys. Rev. A* **53**, R648–51 (1996).
- [49] Z.-T. Lu, C. J. Bowers, S. J. Freedman, B. K. Fujikawa, J. L. Mortara, S.-Q. Shang, K. Coulter, and L. Young, “Laser trapping of short-lived radioactive isotopes,” *Phys. Rev. Lett.* **72**, 3791–4 (1994).
- [50] T. E. Barrett, S. W. Dapore-Schwartz, M. D. Ray, and G. P. Lafyatis, “Slowing atoms with σ^- polarized light,” *Phys. Rev. Lett.* **67**, 3483–6 (1991).
- [51] M. Zhu, C. W. Oates, and J. L. Hall, “Continuous high-flux monovelocity atomic beam based on a broadband laser-cooling technique,” *Phys. Rev. Lett.* **67**, 46–9 (1991).
- [52] W. Ketterle, A. Martin, M. A. Joffe, and D. E. Pritchard, “Slowing and cooling atoms in isotropic laser light,” *Phys. Rev. Lett.* **69**, 2483–6 (1992).
- [53] C. Monroe, Ph.D. thesis, Univ. of Colorado, Boulder, CO 80309, 1993.
- [54] J. A. Behr *et al.*, “Magneto-optic trapping of beta-decaying ^{38}K and ^{37}K from an on-line isotope separator,” *Phys. Rev. Lett.* **79**, 375–8 (1997).
- [55] J. E. Simsarian, A. Ghosh, G. Gwinner, L. A. Orozco, G. D. Sprouse, and P. A. Voytas, “Magneto-optic trapping of ^{210}Fr ,” *Phys. Rev. Lett.* **76**, 3522–5 (1996),
L. A. Orozco, SUNY Stony Brook, private communication.

- [56] V. A. Dzuba, V. V. Flambaum, and O. P. Sushkov, “Calculation of energy levels, E1 transition amplitudes and parity violation in francium,” *Phys. Rev. A* **51**, 3454–61 (1995).
- [57] P. G. H. Sandars, “Enhancement Factor for the Electric Dipole Moment of the Valence Electron in an alkali atom,” *Phys. Lett.* **22**, 290–1 (1966).
- [58] R. Guckert, E. P. Chamberlin, D. W. Preston, V. D. Sandberg, D. Tupa, D. J. Vieira, H. Wollnik, and X. X. Zhao, “Coupling an optical trap to a mass separator,” *Nucl. Instrum. Methods Phys. Res. B* **126**, 383 – 5 (1997).
- [59] L. A. Bray, (Pacific Northwest National Labs, 1996), private communication.
- [60] S. Gerstenkorn, J. Berges, and J. Chevillard, “Atlas Du Spectre D’Absorption de la Molecule d’Iode,” Technical report, Laboratoire Aime Cotton (1982) .
- [61] J. Hall and S. A. Lee, “Interferometric Real-time Display of CW Dye Laser Wavelength with Sub-Doppler Accuracy,” *Appl. Phys. Lett.* **29**, 367 (1976).
- [62] I. collaboration: A. Coc *et al.*, “Isotope shifts, spins and hyperfine structures of $^{118,146}\text{Cs}$ and of some francium isotopes,” *Nuc. Phys. A* **468**, 1 – 10 (1987).
- [63] S. V. Andreev, V. I. Mishin, and V. S. Letokhov, “Rydberg levels and ionization potential of francium measured by laser-resonance ionisation in a hot cavity,” *J. Opt. Soc. Am. B* **5**, 2190 – 8 (1988).
- [64] M. A. Kasevich, E. Riis, L. Chu, and R. G. DeVoe, “RF spectroscopy in an atomic fountain,” *Phys. Rev. Lett.* **63**, 612 – 5 (1989).
- [65] T. Walker, P. Feng, D. Hoffman, and R. S. Williamson III, “Spin-polarized spontaneous-force atom trap,” *Phys. Rev. Lett.* **69**, 2168–71 (1992).
- [66] J. R. Gardner, R. A. Cline, J. D. Miller, D. J. Heinzen, H. M. J. M. Boesten, and B. J. Verhaar, “Collisions of doubly spin-polarized, ultracold ^{85}Rb atoms,” *Phys. Rev. Lett.* **74**, 3764–7 (1995).
- [67] W. Ketterle and N. J. VanDruten, “Evaporative cooling of trapped atoms,” *Adv. At. Mol. Opt. Phys.* **37**, 181–236 (1996).
- [68] G. R. Harrison, Massachusetts Institute of Technology Wavelength Tables (MIT Press, Cambridge, Mass., 1939).
- [69] D. Boiron, A. Michaud, J. M. Fournier, L. Simard, M. Sprenger, G. Grynberg, and C. Salomon, “Cold and dense cesium clouds in far-detuned dipole traps,” *Phys. Rev. A* **57**, R4106–9 (1998).
- [70] K. L. Corwin, Z.-T. Lu, C. F. Hand, R. J. Epstein, and C. E. Wieman, “Frequency-stabilized diode laser with the Zeeman shift in an atomic vapor,” *Appl. Opt.* **37**, 3295–8 (1998).

- [71] M. E. Gehm, K. M. O'Hara, T. A. Savard, and J. E. Thomas, "Dynamics of noise-induced heating in atom traps," *Phys. Rev. A* **58**, 3914–21 (1998).
- [72] Siegman, *Lasers* (University Science Books, Mill Valley, CA, 1986).
- [73] E. Hecht, *Optics* (Addison-Wesley Publishing Company, Reading, Massachusetts, 1990).
- [74] C. G. Townsend, N. T. Edwards, C. J. Cooper, K. P. Zetie, C. J. Foot, A. M. Steane, P. Szriftgiser, H. Perrin, and J. Dalibard, "Phase-space density in the magneto-optical trap," *Phys. Rev. A* **52**, 1423–40 (1995).
- [75] T. P. Dinneen, K. R. Vogel, E. Arimondo, J. L. Hall, and A. Gallagher, "Cold collisions of Sr^* -Sr in a magneto-optical trap," *Phys. Rev. A* **59**, 1216–22 (1998).
- [76] C. S. Adams, H. J. Lee, N. Davidson, M. Kasevich, and S. Chu, "Evaporative cooling in a crossed dipole trap," *Phys. Rev. Lett.* **74**, 3577–80 (1995).
- [77] T. Takekoshi and R. J. Knize, "CO₂ laser trap for cesium atoms," *Opt. Lett.* **21**, 77–9 (1996).
- [78] T. Walker and P. Feng, "Measurements of Collisions Between Laser-cooled Atoms," *Advances in Atomic Molecular and Optical Physics* **34**, 125–70 (1994).
- [79] K. M. O'Hara, S. R. Granade, M. E. Gehm, T. A. Savard, S. Bali, C. Freed, and J. E. Thomas, "Ultrastable CO₂ Laser Trapping of Lithium Fermions," *Phys. Rev. Lett.* **82**, 4204–7 (1999).
- [80] J. D. Miller, Ph.D. thesis, University of Texas, 1994.
- [81] A. Gallagher and D. E. Pritchard, "Exoergic Collisions of Cold Na^* -Na," *Phys. Rev. Lett.* **63**, 957–60 (1989).
- [82] P. D. Lett, K. Mølmer, S. D. Gensemer, K. Y. N. Tan, A. Kumarakrishnan, C. D. Wallace, and P. L. Gould, "Hyperfine structure modifications of collisional losses from light-force atom traps," *J Phys. B* **28**, 65 – 81 (1995).
- [83] P. S. Julienne and J. Vigue, "Cold collisions of ground- and excited-state alkali-metal atoms," *Phys. Rev. A* **44**, 4464–85 (1991).
- [84] S. D. Gensemer, V. Sanchez-Villicana, K. Y. N. Tan, T. T. Grove, and P. L. Gould, "Trap-loss collisions of ^{85}Rb and ^{87}Rb : Dependence on trap parameters," *Phys. Rev. A* **56**, 4055–63 (1997).
- [85] D. Sesko, T. Walker, C. Monroe, A. Gallagher, and C. Wieman, "Collisional Losses from a Light-Force Atom Trap," *Phys. Rev. Lett.* **63**, 961–4 (1989).
- [86] C. D. Wallace, T. P. Dinneen, K.-Y. N. Tan, T. T. Grove, and P. L. Gould, "Isotopic difference in trap loss collisions of laser cooled rubidium atoms," *Phys. Rev. Lett.* **69**, 897 – 900 (1992).

- [87] L. Marcassa, V. Bagnato, Y. Wang, C. Tsao, J. Weiner, O. Dulieu, Y. B. Band, and P. S. Julienne, “Collisional loss rate in a magneto-optical trap for sodium atoms: Light-intensity dependence,” *Phys. Rev. A* **47**, R4563–6 (1993).
- [88] M. B. Peters, D. Hoffmann, J. D. Tobiason, and T. Walker, “Laser-induced ultracold $\text{Rb}(5S_{1/2}) + \text{Rb}(5P_{1/2})$ collisions,” *Phys. Rev. A* **50**, R906–9 (1994).
- [89] D. Hoffmann, P. Feng, and T. Walker, “Measurements of Rb trap-loss collision spectra,” *J. Opt. Soc. Am. B* **11**, 712 – 20 (1994).
- [90] M. Drewsen, P. Laurent, A. Nadir, G. Santarelli, A. Clairon, Y. Castin, D. Grison, and C. Salomon, “Investigation of sub-Doppler cooling effects in a cesium magneto-optical trap,” *App. Phys. B* **59**, 283–98 (1994).
- [91] A. Kuhn, H. Perrin, W. Hänsel, and C. Salomon, “Three Dimensional Raman Cooling using Velocity Selective Rapid Adiabatic Passage,” In OSA Tops on Ultracold Atoms and BEC, K. Burnett, ed., **7**, 58–65 (1996).
- [92] H. Perrin, A. Kuhn, I. Bouchoule, and C. Salomon, “Sideband cooling of neutral atoms in a far-detuned optical lattice,” *Europhys. Lett.* **42**, 395–400 (1998).
- [93] V. Vuletić, C. Chin, A. J. Kerman, and S. Chu, “Degenerate Raman Sideband Cooling of Trapped Cesium Atoms at Very High Atomic Densities,” *Phys. Rev. Lett.* **81**, 5768 – 71 (1998).
- [94] Y. B. Ovchinnikov, D. V. Laryushin, V. I. Balykin, and V. S. Letokhov, “Cooling of atoms on reflection from a surface light wave,” *JETP Lett.* **62**, 113 – 8 (1995).
- [95] P. Desbiolles, M. Arndt, P. Szriftgiser, and J. Dalibard, “Elementary Sisyphus process close to a dielectric surface,” *Phys. Rev. A* **54**, 4292 – 8 (1996).
- [96] C. S. Wood, Ph.D. thesis, Univ. of Colorado, Boulder, CO 80309, 1996.
- [97] J. C. Camparo and R. P. Frueholz, “Parameters of adiabatic rapid passage in the 0-0 hyperfine transition of ^{87}Rb ,” *Phys. Rev. A* **30**, 803–11 (1984).
- [98] M.-O. Mewes, M. R. Andrews, D. M. Kurn, D. S. Durfee, C. G. Townsend, and W. Ketterle, “Output Coupler for Bose-Einstein Condensed Atoms,” *Phys. Rev. Lett.* **78**, 582–5 (1997).
- [99] L. D. Landau and E. M. Lifshitz, Mechanics (Pergamon Press, 1976).
- [100] A. C. Wilson, J. C. Sharpe, C. R. McKenzie, P. J. Manson, and D. M. Warrington, “Narrow-linewidth master-oscillator power amplifier based on a semiconductor tapered amplifier,” *Applied Optics* **37**, 4871–5 (19).
- [101] G. Ferrari, M.-O. Mewes, F. Schreck, and C. Salomon, “High-power multiple-frequency narrow-linewidth laser source based on a semiconductor tapered amplifier,” *Optics Letters* **24**, 151–3 (1999).

- [102] L. S. Cutler, “Frequency stabilized laser system,” U.S. patent 3,534,292 (1970).
- [103] C. E. Wieman and L. Hollberg, “Using diode lasers for atomic physics,” *Rev. Sci. Instrum.* **62**, 1 – 20 (1991).
- [104] K. B. MacAdam, A. Steinbach, and C. Wieman, “A narrow-band tunable diode laser system with grating feedback and a saturated absorption spectrometer for Cs and Rb,” *Am. J. Phys.* **60**, 1098 – 111 (1992).
- [105] B. Cheron, H. Gilles, J. Havel, O. Moreau, and H. Sorel, “Laser frequency stabilization using Zeeman effect,” *J. Phys. III* **4**, 401 – 6 (1994).
- [106] W. Demtröder, Laser Spectroscopy (Springer-Verlag, New York, 1996).