

# **Using ultrafast extreme-ultraviolet pulses for time-resolved dynamics of molecules chemisorbed on metal surfaces**

by  
Chi-Fong Lei

A dissertation submitted in partial fulfillment  
of the requirements for the degree of  
Doctor of Philosophy  
(Electrical Engineering)  
in The University of Michigan  
2003

Doctoral Committee:

Professor Henry C. Kapteyn, Chair  
Professor John Gland  
Professor Margaret M. Murnane  
Professor Stephen Rand

© Chi-Fong Lei 2003  
All Rights Reserved

For Ruth, Matthew and Jacob

## ACKNOWLEDGEMENTS

I would like to thank Profs. Henry Kapteyn and Margaret Murnane for their guidance and patience during my six years of graduate study. They are excellent advisors and have shown me how to tackle problems that I have never seen before, the definition of scientific research. Their insight and knowledge at scientific problems have opened my eyes and led me to the vast world of physics. I especially thank them for giving me opportunities to try many different things, and providing good advices when needed. I also appreciate that they have given me freedom to apply my ideas to the experiments, never forcing me to do something that I did not want to. I thank Henry for his sometimes “nit-picky” requirements – requiring me to not only get the design working, but to also design it cheaply. I think this is good training for my future research – it is important not only to get the experiment done, but to do it well.

I thank Dr. Michael Bauer for his guidance and for bringing me to the field of surface physics. We had many good times doing these experiments, even while staying up very late in the laboratory. Not only did he have brilliant ideas regarding the experiments, he was also an excellent teacher. I would not have come this far if I had not working with him all these years. Furthermore, thanks for his encouragements which kept me going forwards when the experiments were not running well.

Thanks to Dr. Kendall Read for her valuable discussions about the experiments. Even though we worked together for just a couple of months, I appreciate her patience in explaining the fine details to me, when I was very junior and have absolutely no idea what I

was getting into.

I also want to thank many folks in the laboratory. Many thanks to Dr. Sterling Backus, Ron Tobey, and Dr. Lino Misoguti for their help on the laser systems, without their help these experiments would have been impossible. I would also like to thank Prof. John Gland, Prof. Steven Rand, Prof. Dan Dassau, Dr. Yi-De Chuang, Emily Gibson, Dr. Kyoungsik Kim, Dr. Robert Shelton, Dr. Randy Bartels, Prof. Zenghu Chang, Prof. Charles Durfee, Dr. Eric Zeek, David Samuels, and Dr. Erez Gershgoren for their valuable discussions. Thanks to Dr. Yanwie Lui and Prof. David Attwood for making the high quality multilayer mirrors for my setup. And also thanks to the people in the JILA machine shop and electronics shop for their excellent support and equipment making, including Seth Wieman, Hans Green, Blaine Horner, David Alchenberger, Todd Asnicar, Paul Beckingham, Michael Whitmore, and James Fung-A-Fat. For the readability of this thesis, thanks are due to David Samuels, Racheal Tearle, and Marilee DeGoede, who corrected my grammar and writing style.

Finally, I would like to pay the greatest tribute to my family. My wife, Ruth, has sacrificed very much to support me through this entire process, and very likely I cannot ever hope to compensate her for her sacrifices. Thanks also to my mother, and my father- and mother-in-law for their support, both mentally and financially. Finally, my faith in Christianity has been vital to me, enabling me to survive some of the most difficult moments during these years.

# TABLE OF CONTENTS

<b>DEDICATION</b> . . . . .	<b>ii</b>
<b>ACKNOWLEDGEMENTS</b> . . . . .	<b>iii</b>
<b>LIST OF TABLES</b> . . . . .	<b>viii</b>
<b>LIST OF FIGURES</b> . . . . .	<b>ix</b>
<b>CHAPTER</b>	
<b>I. Introduction</b> . . . . .	<b>1</b>
1.1 Desorption induced by intense ultrafast pulses . . . . .	5
1.2 Surface chemistry induced by femtosecond pulses . . . . .	8
1.3 Two-photon photoemission spectroscopy . . . . .	10
1.3.1 Image-potential states . . . . .	11
1.3.2 Electron inelastic decay dynamics on an adsorbate covered surface . . . . .	15
1.3.3 Atomic motion studied by 2PPE . . . . .	19
1.4 Direct measurement of hot surface electrons induced by ultrafast pulses . . . . .	21
1.5 Ultrafast optical methods for surface dynamics investigation . . . . .	24
1.5.1 Probing desorption dynamics with surface second harmonic generation . . . . .	24
1.5.2 Time-resolved vibrational sum-frequency generation . . . . .	26
1.6 Conclusion . . . . .	28
<b>II. Time-Resolved Extreme-Ultraviolet Photoemission Spectroscopy</b> . . . . .	<b>30</b>
2.1 High Harmonic Generation . . . . .	31
2.2 Time-resolved extreme ultraviolet photoemission spectrometer . . . . .	39
2.2.1 Pump - HHG Probe beam line . . . . .	39
2.2.2 Ultra-high Vacuum Chamber . . . . .	44
2.2.3 Finding temporal and spatial overlap . . . . .	50
2.3 Conclusion . . . . .	52

### III. Probing the transition dynamics of chemisorbed states of O<sub>2</sub> on Pt(111) 53

3.1	O <sub>2</sub> states on Pt(111) surfaces . . . . .	53
3.1.1	Physisorbed State . . . . .	53
3.1.2	Superoxo and peroxy chemisorbed states . . . . .	54
3.1.3	Atomic Oxygen . . . . .	58
3.2	Experimental results of O <sub>2</sub> /Pt (111) . . . . .	58
3.2.1	Thermal Desorption Spectroscopy . . . . .	58
3.2.2	Static photoemission spectra of the different oxygen states . . . . .	60
3.2.3	Oxygen desorption induced by intense ultrafast pulses . . . . .	63
3.2.4	Probing O <sub>2</sub> /Pt(111) by TR-UPS . . . . .	65
3.2.5	Permanent modification of O <sub>2</sub> on Pt(111) . . . . .	70
3.3	Discussion . . . . .	71
3.3.1	Nonthermal surface electrons on Pt(111) surface . . . . .	72
3.3.1.1	Nonthermal surface electrons observed in the experiment . . . . .	72
3.3.1.2	Two temperature model and nonthermal surface electrons lifetime . . . . .	73
3.3.1.3	Excitation of O <sub>2</sub> by nonthermal surface electrons . . . . .	78
3.3.2	Mechanism of O <sub>2</sub> excitation . . . . .	80
3.3.2.1	Identification of the transient feature . . . . .	81
3.3.2.2	Excitation rate of molecular oxygen . . . . .	84
3.3.2.3	Energy transfer mechanism from nonthermal electrons to adsorbates . . . . .	85
3.3.2.4	Possible molecular motion during state transition . . . . .	87
3.3.2.5	Decay channels for the excited state . . . . .	88
3.4	Conclusion . . . . .	89

### IV. Probing adsorbate oscillation on metal surfaces . . . . . 91

4.1	Photoelectric current affects by molecular symmetry and EUV polarization . . . . .	92
4.2	Symmetry of the orbitals of CO/Pt(111) . . . . .	96
4.3	Resolving oscillations of adsorbed CO molecules . . . . .	100
4.4	Discussion . . . . .	104
4.4.1	CO normal modes on a platinum surface . . . . .	104
4.4.2	Hot electrons mediated excitation . . . . .	110
4.5	Conclusion . . . . .	113

### V. Future work . . . . . 114

5.1	Time-resolved core-level spectroscopy . . . . .	115
5.2	Probing bimolecular chemical reaction on surfaces . . . . .	116
5.3	Angle- and time-resolved photoemission spectroscopy . . . . .	118
<b>APPENDICES</b>	. . . . .	<b>122</b>
A.1	Desorption induced by electronic transition (DIET) . . . . .	123
	A.1.1 Resonant excitation by non-thermal surface electrons .	125
	A.1.2 Desorption probability . . . . .	130
A.2	Desorption induced by multiple electronic transition (DIMET) .	133
<b>BIBLIOGRAPHY</b>	. . . . .	<b>135</b>

## LIST OF TABLES

### Table

3.1	Comparisons of the adsorption energies and bond lengths between gas-phase oxygen molecules ( $O_2$ ), superoxo ( $O_2^-$ ), peroxy molecular oxygen ( $O_2^{2-}$ ), and atomic oxygen (O) on fcc and hcp sites. . . . .	56
3.2	Parameters of platinum(111) used in the calculation of electron and lattice temperatures . . . . .	75
4.1	Character table of the symmetry group $C_{\infty v}$ . ( $z$ is the direction of the principle symmetric axis) . . . . .	95
4.2	Table of the direct product ( $\otimes$ ) of the symmetry group $C_{\infty v}$ . . . . .	96
4.3	Normal mode energies of CO adsorbed on Pt(111) (* calculated for CO/Ni)	108

## LIST OF FIGURES

### Figure

- 1.1 Desorption yield of two-pulse auto-correlation for NO on Pd(111) surface as a function of the time delay between the two ultrafast pulses. The laser fluences of the two pulses are varied as (a) 1.7 and 1.8 mJ/cm<sup>2</sup>, (b) 1.6 and 2.2 mJ/cm<sup>2</sup>, and (c) 1.3 and 3.0 mJ/cm<sup>2</sup>. (d)-(f) Calculated temperatures of the surface electrons and the lattice phonons. (figure reproduced from Ref. [10]) . . . . . 7
- 1.2 Illustration of the CO/O/Ru(0001) reaction. (a) CO has a much lower desorption barrier comparing to oxygen, which prohibit CO oxidation with the O atom when excited thermally. (b) Under femtosecond pulse excitation, the 1.8 eV barrier for O activation is overcome by coupling from the hot electrons, and CO<sub>2</sub> can be formed. (figure reproduced from Ref. [16]) . . . . . 10
- 1.3 (a) Schematic energy diagram illustrating the process of two-photon photoemission. (b) Image-potential in front of a metal surface. The figure shows the wavefunction probabilities and the energy levels of the  $n = 1$  and 2 image-potential states. The figure also shows the energy band gap of the metal surface, prohibiting the image-potential state electron from penetrating back into the metal surface. (figure reproduced from Ref. [17]) 12
- 1.4 Two-Photo photoemission spectra showing the the image-potential state of an Ag(100) surface at various time delays. The signal is maximized at 0 fs time delay, and gradually decreases with positive time delay. Inset ( $\times 100$  vertical scale): the background spectra at +200 fs time delay due to 2PPE by the single beam excitation (both pump and probe). (figure reproduced from Ref. [19]) . . . . . 14

1.5	(a) Comparison of the full-width-half-maximum inelastic decay times between a Cs covered (0.8 ML) Cu(111) surface and a clean Cu(111) surface shows a remarkable time differences around the cesium unoccupied state between the Fermi edge and the vacuum level of Cu. Data are taken by time-resolved 2PPE. (b) 2PPE spectra of a Cs covered and a clean Cu(111) surfaces clearly shows a Cs induced peak appears at 5.8 eV. The pronounced peak at 6.2 eV of the clean surface (dashed spectrum) arises from the occupied Shockley surface state. (figure reproduced from Ref. [22]) . . . . .	16
1.6	Projected surface band structure for (a) Cu(100) and (b) Cu(111). The dashed line indicates an energy level of an excited alkali state at 2.6 eV above $E_f$ . The momentum distance to real bulk states from the band gap center is much smaller for Cu(100) than for Cu(111). (figure reproduced from Ref. [22]) . . . . .	17
1.7	(a) The 2PPE difference spectra give information on the dynamic surface electronic structure. The 2PPE signal corresponds to one-photon absorption from both the pump and the probe beams, with the delay $\Delta$ set to an integral number of optical cycles. Dashed lines and solid vertical line indicate the Gaussian peak deconvolution of $SS$ and $A$ , and energy of $A$ at $\Delta = 0$ fs, respectively. (b) $E_A(\Delta)$ from two independent experiments for $\theta_{Cs} \sim 0.09$ ML. (c) The wave packet position $R_{Cu-Cs}(\Delta)$ and its polynomial fit. (d) Dashed and solid lines, respectively, represent the predicted and derived excited state PES near the ground-state equilibrium geometry, and the dotted line represents the calculated anti-bonding state energy dependence on $R_{Cu-Cs}$ . (figure reproduced from Ref. [27]) . . . . .	20
1.8	Electron energy distribution function versus energy (reference to the Fermi-edge) taken with the gold film excited by (a) $120 \mu\text{J}/\text{cm}^2$ and (b) $300 \mu\text{J}/\text{cm}^2$ absorbed laser fluence at various time delays. The dashed line is the best Fermi-Dirac distribution fit with the corresponding electron temperatures are shown above the figures. All the figures are drawn in log scale to better shown the hot electrons distribution above the Fermi-edge. (figure reproduced from Ref. [30]) . . . . .	22

1.9	(a) Second harmonic signal versus CO exposure where the Cu(111) surface is under an ambient pressure of $P_{\text{CO}} = 5 \times 10^{-8}$ torr. A single laser shot with fluence of $4.5 \text{ mJ/cm}^2$ to desorb CO molecules from the copper surface. Notice that the second harmonic signal decrease when the surface has higher CO coverage. (b) Second harmonic probe signal versus time delay after CO desorption induced by an ultrafast pump pulse with fluence $F = 4.5 \text{ mJ/cm}^2$ . The autocorrelation of the pump pulse is also shown in the figure. Inset: Same data (solid line) on expanded time scale, along with two other fluences. (figure reproduced from Ref. [34])	25
1.10	(a) SFG spectrum of the C-O stretching vibration of CO/Ru(111) at 340 K and the spectrum of the 150 fs broadband IR pulse (dashed line). Left inset: illustration of time-resolved vibrational sum-frequency generation. (see text) Right side: free induction decay of this vibration exhibiting exponential decay with $\tau = 580 \text{ fs}$ , indicating a homogeneous linewidth of $9.2 \text{ cm}^{-1}$ . (figure taken from Ref. [35]) (b) Transient SFG spectra of the C-O stretch vibration after excitation with 800 nm/100 fs pump pulses with a fluence of $55 \text{ J/m}^2$ , starting at a temperature of 340 K and coverage of 0.33 ML. (figure reproduced from Ref. [36]) . . . . .	27
2.1	High harmonics generated in neon gas at 40 torr with a 1 ps duration, $1.053 \text{ }\mu\text{m}$ Nd laser pulse at an intensity of $1.5 \times 10^{15} \text{ W/cm}^2$ . (figure reproduced from Ref. [50]) . . . . .	32
2.2	In the semi-classical picture, (a) the atomic coulomb potential is disturbed by the laser electric field, and the electron wave packet tunnels through the weaken potential barrier to escape. (b) The electron wave packet enters the free space “continuum” and continue to accelerate in the laser field. On the subsequent half cycle, the electron is decelerated and turns around to be accelerated towards the parent ion. The electron with high kinetic energy then scatters with the parent ion to generate high harmonics. . . . .	34

2.3	Schematic diagram of the hollow-core fiber used to generate the 27 <sup>th</sup> harmonic (42 eV) probe pulse in this experiment. The driving ultrafast beam is drawn in light color and the generated high harmonic beam is in dark color. The fiber has four sections, and the long one is the main harmonic generation region. The noble gas is introduced to the capillary through the gap between the first and second, second and third sections (from the left). The design of having the first and the third short sections sandwiched between the second long section is to maintain a constant pressure at the second section where the high harmonic generation happens. The fourth short section is optionally installed for differential pumping purpose. The inner diameter for all the fibers is 150 $\mu\text{m}$ . The capillary length of all the short sections is 1 cm, and the length for the long section is 8 cm. . . . .	37
2.4	Schematic drawing of the time-resolved ultraviolet photoemission spectroscope. The pump beam is the beam line drawn at the lower part of the figure; the probe beam is drawn at the upper part of the figure, and it goes through the hollow-core capillary to generate high harmonics. The main chamber is under ultra-high vacuum condition at all times ( $\sim 10^{-10}$ torr), and a gate valve is used to separate the main chamber from the rest of the system. The multi-layer mirror chamber and the beam line are under high vacuum ( $\sim 10^{-6}$ torr) condition. . . . .	40
2.5	Combined reflectivity of the Si/Mo multi-layer mirror pair (2 mirrors) used in the TR-UPS setup. The reflectivity of the mirrors are measured by the synchrotron radiation facility in Lawrence Berkeley laboratory. . . . .	43
2.6	The image of the 27 <sup>th</sup> harmonic fluoresce at a phosphor screen. The phosphor screen is mounted beneath the platinum sample, and the phosphor screen was raised to the sample position when this picture was taken. The image was capture by a low-lux, high sensitive camera mounted outside the UHV chamber and looking in through a view port. The spot size of the harmonic is measured to be 400 $\mu\text{m}$ . The spot seems elliptical because the phosphor screen is at 45 degrees with respect to the incoming EUV beam. . . . .	43
2.7	The output signal of the impedance matched MCP TOF detector when a single electron hits the detector. The signal is directly measured from the detector with an oscilloscope. It is very important that there is no signal reflection or “ringing” after the main peak for high count rate electron detection. . . . .	47

2.8	Picture shows the internal arrangement of the TR-UPS chamber. In this photo, the sample is turned 90 degrees from the normal position for better viewing. Both of the pump-probe beams are coming in from the flange at the left side of the picture to reach the metal sample. . . . .	49
2.9	Signal of second harmonic generation between the infrared pump and probe pulses. The FWHM of the signal is 62 fs. . . . .	51
3.1	Schematic diagram of the surface configuration of oxygen on Pt (111), for the superoxo ( $O_2^-$ ), peroxo ( $O_2^{2-}$ ), and atomic (O) oxygen states. . .	54
3.2	TDS for Pt(111) covered with $O_2$ : the rate of temperature increase is 3 K/sec from 77 K to 300 K; then 10 K/sec from 300 K to 1100 K . . . .	59
3.3	Static photoemission spectra taken for the superoxo ( $O_2^-$ ), peroxo ( $O_2^{2-}$ ) and atomic (O) oxygen states on Pt(111). . . . .	62
3.4	Femtosecond laser-induced desorption for the two adjacent desorption cycles with oxygen redosing between the two laser excitations. Insert shows the integrated desorption signal for the two cycles. . . . .	66
3.5	Electron energy distribution at Fermi edge of $O_2$ /Pt(111), for time delays of 0 fs, 150 fs, 250 fs, and 2 ps. Nonequilibrium hot electrons can be seen at 0 and 150 fs; however, at 250 fs, the energy distribution has thermalized. Inset: Fermi edges at 0 fs time delay (black square) and static spectrum without pump pulses excitation (hollow square), plotted in log scale. The electron distribution at 0 fs time delay can be fit to a 100 K Fermi-Dirac distribution with 6% of the electrons excited by the pump photon energy (1.5 eV). . . . .	67
3.6	Time-resolved high-resolution photoemission spectra from a saturation layer of molecular oxygen adsorbed on a Pt(111) surface at liquid nitrogen temperature: (a) no pump beam; (b) with pump beam, zero time delay between pump and probe; (c) 250 fs time delay between pump and probe; (d) 500 fs time delay; (e) repeat of (b), taken immediately after (d). . . . .	68
3.7	Integrated amplitude of the characteristic 6 eV peak feature as a function of pump-probe delay. A fit to the data indicates an onset time of $550 \pm 140$ fs. . . . .	69

3.8	Spectra taken during pump beam irradiation of the sample over a period of 2 hours. During the first $\approx 50$ min, the spectra are unchanged. However, after $\approx 100$ min, the characteristic 6 eV peak gradually increases and is permanently present on the subsequent spectrum taken. . . . .	71
3.9	Calculated electron temperature ( $T_e$ ) and lattice temperature ( $T_p$ ) using the “two-temperature model” at the platinum (111) surface excited by a $10 \mu\text{J}/\text{mm}^2$ , 55 fs ultrafast infrared pulse. The measured nonthermal electron integrated intensity (square) is also shown in the figure for comparison purposes. . . . .	76
3.10	Integrated intensity of the peroxo signature peak compared to the rescaled (a) electron and (b) lattice temperature. . . . .	79
3.11	Because of the increased population of the surface electrons above the Fermi level induced by the ultrafast pump pulse excitation, the surface electrons will start to tunnel to the $1\pi_g^*$ orbital of the oxygen molecule and this leads to energy transfer to the oxygen center-of-mass. . . . .	86
3.12	Possible reaction diagram of how oxygen molecules react after excited by an intense ultrafast pulse on the platinum surface . . . . .	89
4.1	Schematic diagram showing the bonding character of a CO molecule adsorbed on a metal surface. (a) The CO-metal bond is provided by the most extended orbital of the molecule which is the $5\sigma$ orbital. Charge back donation from the metal into the formerly empty $2\pi^*$ orbitals is indicated by thin arrows. (b) Schematic energy level diagram of the CO-metal system. The energy levels on both sides are for the isolated CO molecule and the metal surface. The middle diagram illustrates the energy levels of an adsorbed CO molecule. Band splitting for the adsorbed CO happens for the $5\sigma$ and $2\pi^*$ levels due to hybridization with the metal $d$ -band and the $p$ -orbitals. (figures reproduced from Ref. [92]) . . . . .	97
4.2	Photoemission spectrum of CO molecules adsorbed on a platinum(111) single crystal surface. Spectra are taken with high harmonic photons (42 eV) with different $s$ (black) and $p$ (red) polarizations. . . . .	99

4.3	(a) Schematic diagram illustrating that when the CO molecules are aligned with the polarization of the EUV pulse, the photoemission cross-section is maximized. After excitation by an ultrafast laser pulse, the CO molecules start to vibrate. When the CO molecules slightly non-parallel to the EUV polarization, the photoemission cross-section is reduced. (b) Pump-probe spectra close to the $4\sigma$ peak. These spectra are taken when the polarization of the EUV probe pulses is perpendicular to the platinum surface ( $p$ -polarized), and the polarization of the excitation pump pulses are parallel to the surface ( $s$ -polarized) with $15\mu\text{J}/\text{mm}^2/\text{pulse}$ fluence. Spectra of 0 fs and 400 fs time delays are shown for comparison, showing that the the $4\sigma$ peak in both height and area at 0 fs time delay are bigger than those at 400 fs time delay. . . . .	101
4.4	Integrated area of the $4\sigma$ peak of the CO/Pt(111) plotted at various time delays, showing that the CO molecules are oscillating at the platinum surface after pump pulse excitation. . . . .	103
4.5	Pump-probe spectra near the Fermi-edge, showing that hot electrons are rapidly excited when the ultrafast pump pulses excites the surface. They subsequently relax in about 200 fs. There is a static component resulting from a small residual 29 <sup>th</sup> harmonic. . . . .	105
4.6	Schematic representation of the normal modes of on-top and bridge-bonded CO on platinum surface. The relative length of the arrows represent the movements of the carbon and oxygen atoms. (figure adapted from Ref. [100] and Ref. [101]) . . . . .	106
4.7	(a) When an ultrafast pulse excites the platinum surface, hot electrons are created on the surface. The hot electrons then tunnel to the $2\pi^*$ orbital of the CO molecule. (b) Due to the bonding character to the platinum surface of the $2\pi^*$ orbital, the C-O vibrational mode $\nu_1$ and the CO-Pt vibrational mode $\nu_2$ will be induced. The frustrated translation mode $\nu_4$ could either be directly excited by the hot electron, or subsequently excited due to energy coupling between modes. . . . .	110

5.1	(a) Schematic drawing of “cold-target recoil-ion momentum spectroscopy” (COLTRIMS). A two-stage supersonic gas jet is installed to provide a well-defined atomic beam, which collides with an energetic ion-beam. Helmholtz coils and resistive coated parallel plates are used to generate homogeneous magnetic and electric fields to confine the recoil-ions and the ejected electrons. Position-sensitive micro-channel plates (MCP) are used to resolve the kinetic energy and emission angles of the recoil-ions and electrons. (figure reproduced from Ref. [113]) (b) The anode of the position-sensitive MCP is wrapped by a fine copper-wire (delay-line) on each dimension. When an electron strikes the anode, the position of the electrons on the detector can be extracted by measuring the time difference of the electronic signals at both ends of each delay-line. The position for each dimension can be determined independently. (figure reproduced from Ref. [114]) . . . . .	120
A.1	Energy level diagram showing that the non-thermal electrons are energized rapidly by an intense ultrafast pulse. These “hot” electrons could scatter into the LUMO state of the adsorbates to transfer energy to excite the molecular motions through Franck-Condon excitation scheme. (figure reproduced from Ref. [79]) . . . . .	127
A.2	Potential curves illustrating the process of Desorption Induced by Electronic Transition (DIET). (a) Intra-molecular electronic excitation involving bound-to-repulsive excitation; (b) Intra-molecular electronic excitation involving attractive intermediate state mediation. (figure reproduced from Ref. [79]) . . . . .	129
A.3	Potential energy curves of center-of-mass translational motion of the chemisorbed ion (charge transfer from surface) with respect to the surface, showing wavepacket propagation throughout the time sequence involving the negative-ion LUMO resonance. The distribution of the final adsorbate states, between vibrationally excited bound (white) and desorptive continuum states (black) is shown as $P(\epsilon)$ versus $\epsilon$ . (figure reproduced from Ref. [79]) . . . . .	132
A.4	Trajectories for motion on the ground and excited PES. The dotted lines represent the total energy of the adsorbate complex, with vertical segments arising from Franck-Condon transitions and horizontal segments from conservative motion on a given PES. (a)(b) Trajectories displayed associated with a single-excitation DIET process. (c)(d) Trajectories for the multiple-excitation DIMET process. (figure reproduced from Ref. [11])	134